

## Breeding strategies for hybrid rice parental line improvement

AS Hari Prasad\*, P Senguttuvel, P Revathi, KB Kemparaju, K Sruthi, RM Sundaram, M SeshuMadhav, MS Prasad and GS Laha

ICAR-Indian Institute of Rice Research, Hyderabad, India

\*Corresponding author e-mail: hari.prasad1@icar.gov.in

### ABSTRACT

*In our country, significant progress has been made in the development of hybrid rice parental lines both female and male parents with good floral traits. An array of hybrids with good grain quality (both unique Basmati type and medium slender (MS) grain type) is developed. Efforts are focussed on the development of restorer lines through exploitation of inter sub-specific hybridization, recurrent selection and population improvement that will help in the development of highly heterotic rice hybrids. The female parental lines are being improved for traits like stigma exertion and high outcrossing that helps in higher seed yields, thereby reducing the hybrid seed cost. Efforts are on to develop appropriate parental lines that can help in the development of suitable hybrids in late duration group and also for unfavourable ecologies. Many newly developed parental lines are being fortified with resistance genes for major diseases such as BB, blast and insect pests such as BPH that will help in the development of hybrids having resistance to major pests and diseases.*

**Key words:** Hybrid rice, CMS line, maintainer, restorer lines, gene pools

### INTRODUCTION

Hybrid rice is a proven and viable technology to enhance the rice production and productivity and it was launched in India in 1989, after its successful adaptation in China. As a result of intensive research efforts over the last three decades, a total of 97 rice hybrids were released for commercial cultivation in different rice growing regions of the country. The area planted to hybrid rice in the country during *Kharif* 2017 was around 3m.ha. (5% of the total rice area of 44 m.ha) and has contributed 3-4 m.t. of additional rice to the total rice production in the country.

However, its adoption is rather slow than the expected level. The main reasons for this are marginal heterosis, narrow genetic base of the parental lines, high seed cost because of seed production issues, non-availability of hybrids in medium late/late duration groups besides quality concerns. All these concerns can be addressed through focussed parental line improvement

by developing an array of appropriate genotypes that can help in the development of hybrids with enhanced yield heterosis, better grain quality and resistance to major biotic stresses. In this paper, different approaches and strategies for parental lines improvement is presented.

### Diversification of CMS sources

All the hybrids released in the country are based on a single source of cytoplasmic male sterility (CMS) *viz.*, Wild Abortive (WA) system. Dependence on a single CMS source on a long run may result in genetic vulnerability of hybrids to sudden outbreak of diseases and insect pests. Hence, it is required to develop CMS lines with diverse cyto-sterility sources. Wild species of rice such as *Oryza nivara*, *Oryza rufipogon* and *Oryza perennis* were already found to have alternate cyto-sterility sources, but their commercial exploitation was limited by the fact that no good restorers are available for these CMS sources like WA system. Intensive efforts are being made to identify suitable

restorers for these various sterility sources. Another sterility source Kalinga I and a CMS line CRMS 32A is developed using this source and a hybrid *viz.*, Rajlaxmi has also been developed from this new source (NRRI, Cuttack).

The People's Republic of China (where the hybrid rice technology is originated) has succeeded in diversifying the CMS sources. Though CMS WA-type remains the dominant type in three line hybrid rice production, they could able to reduce the dependency on this system from 69% in 1996 to 47% in 2003. Among the other types of CMS, CMS-ID has been used widely in recent years because of its good flowering habits and grain quality. Its proportion has increased from less than 10% in 1996 to more than 20% in recent years (Shi-hua et al., 2009). Another CMS system *viz.*, CMS-G&D has also been successfully deployed in the development of hybrids in China (Shi-hua et al., 2004).

### Improvement of CMS & Maintainer Lines

One of the major limitations of the hybrid rice technology is the higher seed cost and it's acting as deterrent in wide spread adoption of hybrid rice cultivation in the country. This can be overcome by developing CMS and maintainer lines with good floral traits such as high outcrossing ability etc. To achieve this, single, double and multiple crosses among maintainers and outstanding partial maintainers (having desirable traits) may be used to develop large segregating generations. The population size in  $F_2$  should be sufficiently large ( $> 2000-3000$  plants) and careful selection has to be exercised for outcrossing and combining ability related traits by retaining sufficient genetic diversity of the segregants. Selection for plant type, grain type, stigma exertion (Takano Kai et al., 2011), elongated uppermost internode (Singh et al., 2007) and other easily observable traits can be made in early segregating generations and combining ability of the desirable fixed lines will be tested at  $F_5/F_6$  stage after confirming the presence of stable maintainer genes, through test crossing.

Two promising maintainer lines *viz.*, APMS 6B, IR 68897B were improved for stigma exertion trait through back cross breeding approach with donor genotypes *viz.*, BF 16B and BF 2096 (Singh et al., 2015). Another popular maintainer line IR 58025B has been improved for elongated uppermost internode-eui

(Rahul et al., 2016).

It's essential to improve the maintainer lines for major biotic stresses, so that the same can be incorporated in CMS lines, which will help in the development of hybrids with resistance to major biotic stresses. IR 58025B has been fortified with bacterial blight (BB) and blast resistance genes through marker-assisted backcross breeding strategy (Hari et al., 2013). By adopting the similar strategy, two more maintainer lines *viz.*, Pusa 6B for BB resistance (Basavaraj et al., 2009); DRR 17B for BB, blast, gall midge resistance (Balachiranjeevi et al., 2015) have been improved. Promising maintainers like these, are being converted into new/improved CMS lines through recurrent back crossing.

### Improvement of Restorer Lines

To increase the yield heterosis of the rice hybrids from the present 15-20% level to at least 25-30%, it's essential to focus on the restorer line improvement by adopting different recombination breeding strategies *viz.*, pedigree, backcross, incomplete backcross, single seed decent, modified backcross methods, multiple convergent improvement, genetic male sterility facilitated recurrent selection methods.

Choice of parents depends on the objective of the breeding program. Genetically diverse donors for various traits like better plant type (tropical japonica parents possessing wide compatibility genes), better restoration ability, high pollen production, relatively tall stature, better grain quality as per the local requirements, multiple disease and insect pest resistance, good GCA and SCA etc. should be used in the crossing programme. Single, three way and multiple crosses among restorers and partial restorers may be used to develop large segregating generations. Depending on the purpose the type of crosses may be  $R_1 \times R_2$ ,  $R_1 \times (R_2 \times R_3)$ ,  $R_1 \times PR$ ,  $R_1 \times (PR \times R_2)$ ,  $R_1 \times (PR_1 \times PR_2)$ ,  $PR \times (R_1 \times R_2)$ ,  $PR_1 \times (R_1 \times PR_2)$ ,  $(R_1 \times R_2) \times (R_3 \times R_4)$ ,  $(R_1 \times PR_1) \times (R_2 \times PR_2)$  and so on.

It's essential to improve the restorer lines for major biotic stresses, so that hybrids with resistance can be developed. The popular restorer lines *viz.*, KMR 3 (Hari et al., 2011); PRR 78 (Basavaraj et al., 2010, Lalitha Devi et al., 2013) for BB resistance; PRR 78 for blast resistance (Vikas et al., 2012); RPHR 1005

for BB and blast resistance (Abhilash Kumar et al., 2016) have been improved through marker-assisted backcross breeding strategy. Efforts are on to improve few more restorer lines *viz.*, BCW 56, EPLT 109, DR 714-1-2R, in a similar way.

Genetic male sterility (GMS) facilitated population improvement is carried out for improvement of restorer lines. As against quick fixation of genes during selfing generations of recombination breeding, genetic male sterility facilitated recurrent selections provide opportunities for continuous recombination, accumulation of favorable genes, broadening of the genetic base and breaking of undesirable linkages. By combining this strategy with Marker Assisted Selection (MAS), efforts are on develop a new set of restorer lines having resistance to multiple biotic stresses.

Inter sub-specific (*indica* x tropical *japonica*) hybridization approach, will help in developing parental lines that can be utilized in identification of hybrids with enhanced yield heterosis. However, a major difficulty encountered in the development of such inter-sub specific hybrids is the partial hybrid sterility (HS) frequently observed in most *indica/japonica* crosses. A special class of rice germplasm, known as wide compatible varieties (WCVs), can produce hybrids with normal fertility when crossed with both *indica* and *japonica*. Out of several genes reported to be involved in hybrid sterility, the S5 locus on chromosome 6 is considered to be the major. Sundaram et al., (2010) developed a functional marker which can distinguish all the three allelic states (*i.e.*, *indica*, *japonica* and neutral at S<sub>5</sub>; this marker is being effectively used in hybrid rice breeding programme to identify the lines having S<sub>5</sub> neutral alleles (Revathi et al., 2010).

Certain popular varieties of our country like Swarna, Samba Mahsuri, which are considered as mega varieties occupying substantial area under cultivation across the country and it's desirable if we develop hybrids similar to these mega varieties. However, these varieties can't be used in the hybrid breeding programme for the reason that they are found to be partial restorers. In order to use these genotypes, efforts are being made to convert them into usable restorer lines, by crossing them with popular restorer lines *viz.*, KMR-3, IBL-57 and trying to identify the promising restorers in the advance segregating generations. A set of restorer lines

are developed and identified, following this strategy and they are being utilized in the development of new hybrid combinations.

Wild species of rice are also being used in the improvement of restorer lines. Of the two major effect yield enhancing QTLs *yld2.1* and *yld8.2* mapped from an Indian accession of *Oryza rufipogon*, *yld2.1* was introgressed into KMR 3, a restorer line of popular hybrid KRH 2 and several introgression lines of KMR 3 with upto 20% yield increase have been obtained (Sudhakar et al., 2012) and these restorers are being used in the development of new hybrids.

### Development of heterotic gene pools

A concept, well exploited in corn in development of gene pools that led to development of highly heterotic hybrids, needs to be exploited in the improvement of hybrid rice parental lines. The work has been initiated in this direction by pooling restorer as well as maintainer lines and to study the extent of diversity among the lines and after classification in to different groups, selected lines from each group will be intermated with other lines from each group and also with lines from other groups to estimate their combining ability and heterosis. Our aim is to develop initially gene pools in different duration groups following this strategy.

All these strategies will hopefully lead to the development of better parental lines and hybrids, helping in the expansion of hybrid rice area and increased production and productivity in the country.

### REFERENCE

- Kumar AV, Balachiranjeevi CH., Naik SB, Rambabu R., Rekha G, Madhavi KR, Harika G, Vijay S, Pranathi K, Hajira SK, Srivastava A, Mahadevaswamy HK, Anila M, Yugander A, Aruna J, Hari Prasad AS, Madhav MS, Laha GS, Viraktamath BC, Balachandran SM, Senguttuvel P, Kemparaju B, Ravindra Babu V, Sundaram RM and Prasad MS (2016). Marker-assisted introgression of the major bacterial blight resistance gene, *Xa21* and blast resistance gene, Pi54 into RPHR-1005, the restorer line of the popular rice hybrid, DRRH3. *J. Plant Biochem. Biotechnol.* PP: DOI 10.1007/s13562-016-0352-z
- Singh AK, Kemparaju K. B, Patil V, Priyadarshini R, Koradi P, Khandekar D, Jayaramulu K, Sheshu Madhav M, Lalitha Shanthi M, Khera P, Barthwal M, Subba Rao LV, Hariprasad AS, Ulaganathan V and Vinay

- Shenoy V (2015). Stigma exertion trait in rice (*Oryza Sativa* L.) And comparison of two Phenotyping methods. *International Journal of Current Research*, 7(03): 13123-13135
- Balachiranjeevi CH, Bhaskar Naik S, Abhilash V, Akanksha S, Viraktamath BC, Madhav MS, Hari Prasad AS, Laha GS, Prasad MS, Balachandran SM, Neeraja CN, Satendra Kumar M, Senguttuvel P, Kemparaju KB, Bhadana VP, Ram T, Harika G, Mahadevaswamy HK, HajiraSk, Yugander A, Pranathi K, Anila M, Rekha G, BVN Kousik M, Dilipkumar T, Swapnil RK, Archana Giri and Sundaram RM (2015). Marker-assisted introgression of bacterial blight and blast resistance into DRR17B, an elite, fine-grain type maintainer line of rice. *Molecular breeding* 35:15
- Basavaraj SH, Singh VK, Singh A, Singh D, Nagarajan M, Mohapatra T, Prabhu KV and Singh AK (2009). Marker aided improvement of Pusa 6B, the maintainer parent of rice hybrid Pusa RH10, for resistance to bacterial blight, *Indian J. Genet.* 69 (1): 10-16
- Basavaraj SH, Singh VK, Singh A, Singh A, Singh A, Anand D, Yadav S, Ellur RK, Singh D, Gopalakrishnan S, Nagarajan M, Mohapatra T, Prabhu KV and Singh AK (2010). Marker-assisted improvement of bacterial blight resistance in parental lines of Pusa RH10, a superfine grain aromatic rice hybrid. *Mol. Breed.* 26: 293-305
- Cheng Shi-hua, Cao Li-yong, Yang Shi-hua, Zhai Hu-qu (2004). Forty Years' Development of Hybrid Rice: China's Experience, *Rice Science* 11(5-6): 225-230
- Cheng Shi-hua, Cao Li-yong, ZhaunagJie-Yun, Wu Weiming, Yang Shi-hua and Zhan Xiao-deng (2009). A breeding strategy for hybrid rice in China, In 'Accelerating hybrid rice development' (eds. Xie F, Hardy B), International Rice Research Institute, Philippines pp. 25-34
- Hari Y, Srinivasarao MS, Viraktamath BC, Hari Prasad AS, Laha GS, Ilyas Ahmed M, Natarajkumar P, Sujatha K, Prasad MS, Pandey M, Ramesha MS, Neeraja CN, Balachandran SM, Shobha Rani N, Kemparaju KB, Madhan Mohan K, Sama VSAK, Sk. Hajira, Balachiranjeevi CH, Pranathi K, Reddy GA, Madhav MS and Sundaram RM (2013). Marker-assisted introgression of bacterial blight and blast resistance into IR 58025B, an elite maintainer line of rice, *Plant Breeding* 132, 586-594 (doi:10.1111/pbr.12056)
- Hari Y, Srinivasarao K, Viraktamath BC, Hari Prasad AS, Laha GS, Ahmed MI, Kumar PN, Ramesha MS, Neeraja CN, Balachandran SM, Rani NS, Suresh PB, Sujatha K, Pandey M, Reddy GA, Madhav MS and Sundaram RM (2011). Marker Assisted improvement of a stable restorer line, KMR-3R and its derived hybrid KRH-2 for bacterial blight resistance and grain quality, *Plant Breeding* doi:10.1111/j.1439-0523.2011.01881.x
- Devi GL, Koradi P, Shenoy V and Shanti ML (2013). Marker-assisted selection: evaluation of bacterial blight resistance in the pyramids of fertility restorer lines. *Journal of Plant Pathology* 95 (2): 299-311
- Priyadarshi R, Hari Prasad AS, Singh AK, Ulaganathan K and Shenoy V (2016). Comparative analysis of effectiveness of wide compatibility (wc) trait between improved maintainer line having wca and eui genes and maintainer line with eui and without wca genes in rice (*Oryza Sativa* L.). *Environment & ecology* 34 (4b): 214
- Revathi P, Singh AK, Sundaram RM, Senguttuvel P, Kemparaju KB, Hari Prasad AS and Viraktamath BC (2010). Molecular screening for the presence of wide compatibility gene S5 neutral allele in the parental lines of hybrid rice. *Indian J. Genet.* 70(4): 373-376
- Singh S, Gangashetti MG and Khera P (2007). Development of elongated uppermost internode CMS lines for hybrid rice breeding in India. *Molecular Plant Breeding* 5(2): 215-216
- Singh VK, Singh A, Singh SP, Ellur RK, Choudhary V, Singh D, Gopala Krishnan S, Nagarajan M, Vinod KK, Singh UD, Prashanthi SK, Agrawal PK, Bhatt JC, Mohapatra T, Prabhu KV, Sarkel S, Rathore R and Singh AK (2012). Incorporation of blast resistance into "PRR78", an elite Basmati rice restorer line, through marker assisted backcross breeding. *Field Crops Res* 12:8-16
- Sudhakar T, Batchu AK, Sarla N and Ramanan R (2012). Os11Gsk gene from a wild rice, *Oryza rufipogon* improves yield in rice. *Funct. Integ. Genom.* 12: 277-289
- Sundaram RM, Sakthivel K, Hari Prasad A S, Ramesha MS, Viraktamath BC, Neeraja CN, Balachandran SM, Shobha Rani N, Revathi P, Sandhya P and Hari . (2010). Development of a PCR based functional marker system for the major wide compatible gene locus S<sub>5</sub> in rice. *Mol. Breeding* 26: 719-727
- Takano-Kai N, Doi K and Yoshimura A (2011). GS3 participates in stigma exertion as well as seed length in rice. *Breeding Science* 61: 244-250